



Shock your litter back to health

Litter Shock “shocks” your litter back to health by adding billions of beneficial *Bacillus* bacteria to the litter. Soaked for 24 hours, **Litter Shock's** bacteria fully activate and increase their numbers by up to 4000%. Spraying all of these beneficial bacteria onto the litter provides a biological “shock treatment” that immediately begins to rebalance the litter microbiome and creates a healthier growing environment.

- ✓ Out-competes, and competitively excludes disease-causing bacteria to reduce disease outbreaks.
- ✓ Dries, composts, and improves litter quality to allow more growouts on the same built-up litter.
- ✓ Reduces ammonia production to reduce stress on birds.
- ✓ Reduces mortality, improves weight gain and feed conversion.

How to use Litter Shock:

- ✓ **Jump-start your regular IMPACT-P(NA) litter management program:** Apply at 1-lb./2,000 sq. ft. 1 day before applying IMPACT-P(NA).
- ✓ **Mitigate dermatitis outbreaks:** Apply at 1-lb./1,000 sq. ft. at first signs of an outbreak.
- ✓ **Reduce Salmonella exposure during feed withdrawal:** Apply at 1-lb./1,000 sq. ft. 1-2 days before starting feed withdrawal.
- ✓ **Reduce ammonia levels naturally and safely:** Apply at 1-lb./1,000 sq. ft. at first signs of increasing ammonia.

